

## Kinetic study of oxidation of mixture of Malonic acid + Tartaric acid by Chromic acid in presence of Hydrochloric acid and salts

DEEPTI KHARE\* and A.H. ANSARI\*\*

\*Patel College of Science and Technology, Bhopal (India)

\*\*Saifia Science College Bhopal (India)

Email:-[deepti.khare@yahoo.co.in](mailto:deepti.khare@yahoo.co.in)

(Acceptance Date 26th February, 2016)

### Abstract

The kinetics of oxidation of mixture of Malonic acid +Tartaric acid by Chromic acid has been studied in presence of Hydrochloric acid at 28°C. The reaction follows first order kinetics with respect to each substrate, oxidant and hydrogen ion concentration. Ostwald isolation method is used for study of rate of reaction. With increase in the concentration of Hydrochloric acid the rate constant of the system increased. The oxidation of mixture of acids was studied at different concentration of Hydrochloric acid. The rate constant, catalytic constant, dielectric constant and temperature coefficient have been determined for all the runs. The probable products of oxidation are CO<sub>2</sub> and carbonyl dicarboxylic acid. Various thermodynamic parameters like energy of activation and frequency factor were also determined. The probable mechanism with result obtained has been proposed with the following rate equation.

$$\frac{d[\text{Cr(VI)}(\text{Hydrochloric acid})]}{dt} = k [\text{Cr (VI)}][\text{Malonic acid}][\text{Tartaric acid}][\text{H}^+]$$

*Key words:* Oxidation, Ostwald isolation, dielectric constant, catalytic constant, frequency factor.

### Introduction

Hexavalent Chromium<sup>1-8</sup> has been extensively used in analytical and preparative chemistry. Thus Chromic acid, aqueous

dichromate, Chromyl chloride, Chromyl acetate and ditertiary butyl chromate are important chromium (VI) compounds used in the oxidation of variety of organic and inorganic substrate in solutions. The chemistry and reactivity pattern of these chromium compounds

has been reviewed critically and extensively<sup>3-9</sup>.

The oxidation of Malonic acid by various metal ion oxidants has been reported with different results. The oxidation of Malonic acid and tartaric acid by Chromic acid exhibit a first order reaction kinetics. Sengupta and other coworkers<sup>15-26</sup> did not undertake Hydrogen ion concentration dependence.

## Experimental

### Preparation of reagents :

All chemicals and reagents used in the work were analar grade and were manufactured by Merck, Reidel. Corning glassware's were employed and doubly distilled water is used through out the study, for preparation of solution. The rate of reaction of the oxidant and reductant were studied by monitoring the decrease in optical density using digital colorimeter at 420nm and pH value by digital pH meter

Systronic model 335(electronic) also refractive index is measured by Abbe's refractometer. All kinetic measurements were carried out and CO<sub>2</sub> and Carbonyl dicarboxylic acid has been found as end products. These products were identified by Physico chemical methods<sup>18</sup>. And they are confirmed by their usual tests.<sup>10-14</sup>

### Product Analysis :

Product analysis was carried out under kinetic conditions. The studies were carried out in aqueous medium and at three different temperatures range of 28°C, 38°C and 48°C. All the solutions were kept in a Toshniwal thermostat at constant temperature with accuracy  $\pm 0.10^\circ\text{C}$ . The required volume of these solutions for each run were mixed and analysed by colorimeter.

The first order reaction was obtained, and the different measured values are shown on the following tables:

Rate constants, pH values, H<sup>+</sup> ion concentration and K/H<sup>+</sup> values

Table 1

S.No.	Concentration of Hydrochloric Acid (M)	Rate Constants $K \times 10^{-2} (\text{min}^{-1})$	pH at 28°C	[H <sup>+</sup> ] ion Concentration $\times 10^{-2}$	K/H <sup>+</sup>
1	0.0000	2.89	2.32	0.48	-
2	0.1000	3.18	2.28	0.52	6.115*
3	0.2000	3.48	2.22	0.60	5.800
4	0.3000	3.65	2.15	0.71	5.141
5	0.4000	3.89	2.13	0.74	5.257
6	0.5000	4.10	2.09	0.81	5.062
7	0.6000	4.25	2.07	0.85	5.000


### Salt Effect

In present investigation the author has observed the effect by increasing concentration of salts on the reaction mixture and retarding effect is observed on the rate constant.

Table 2


S. No.	Concentration of salts(M)	$K \times 10^{-2} (\text{Min}^{-1})$			
		NaCl	KCl	$\text{Na}_2\text{SO}_4$	$\text{K}_2\text{SO}_4$
1	0.0000	2.89	2.89	2.89	2.89
2	0.1000	2.64	2.48	2.71	2.59
3	0.2000	2.41	2.32	2.56	2.34
4	0.3000	2.34	2.10	2.37	2.10
5	0.4000	2.10	2.00	2.12	2.04
6	0.5000	1.87	1.74	2.01	1.91
7	0.6000	1.78	1.64	1.88	1.74

### Variation of rate constant with concentration of Hydrochloric acid



Cf. Table no. 1

### Variation of rate constant with concentration of salts



Cf. Table no. 2

*Effect of Temperature :*

The experiment was done at three different temperatures *ie.* 28°C, 38°C and 48°C. and observation was that the rate of the reaction become almost doubled at every 10°C rise in temperature. Hence reaction is normal and homogeneous and adhere to Arrhenius equation.

**Rate constant at three different temperatures and pH values**

Table 3

S.No.	Concentration of Hydrochloric Acid (M)	Rate Constants $K \times 10^{-2} \text{min}^{-1}$			pH at
		28°C	38°C	48°C	28°C
1	0.1000	3.18	6.23	12.65	3.18
2	0.2000	3.48	6.99	13.92	3.48
3	0.3000	3.65	7.26	14.60	3.65
4	0.4000	3.89	7.66	15.63	3.89
5	0.5000	4.02	8.12	16.08	4.10
6	0.6000	4.25	8.33	16.74	4.25

**Temperature coefficients for all the reaction mixtures**

Table 4

S.No.	Concentration of Hydrochloric Acid (M)	Temperature coefficient	
		$K_{38}/K_{28}$	$K_{48}/K_{38}$
1	0.1000	1.96	2.03
2	0.2000	2.01	1.99
3	0.3000	1.99	2.01
4	0.4000	1.97	2.04
5	0.5000	2.02	1.98
6	0.6000	1.96	2.01

**Catalytic constant and dielectric constant**

Table 5

S.No.	Concentration of Hydrochloric Acid (M)	Catalytic Constant $\times 10^{-1}$	Dielectric Constant
1	0.1000	05.58*	1.686
2	0.2000	09.83	1.654
3	0.3000	10.70	1.639
4	0.4000	13.51	1.631
5	0.5000	14.94	1.629
6	0.6000	16.00	1.612
	Average	13.00	1.642

\*Not included in average.

## Result and Discussion

The oxidation of mixture of Malonic acid + Tartaric acid by Chromic acid were kinetically studied. In our investigation three main factors which affect the order of a chemical reaction, these are

- 1) Substrates
- 2) Oxidant
- 3) Hydrogen ion (H<sup>+</sup>)

Here in this investigation order of reaction is one with respect to substrate, oxidant and [H<sup>+</sup>], hence rate of reaction is affected by the extent of acid which depends on certain factors like;

- Nature of added acid
- Concentration of added acid

In this reaction it is observed that the pH value of the reaction mixture do not change much by increasing concentration of Hydrochloric acid.

The average values of Catalytic constant is found to be  $13 \times 10^{-1}$

Energy of activation  $\Delta E^\ddagger = 12.754$  KCals and Frequency factor  $A = 7.339 \times 10^7$

Hence the rate law is given as

$$-\frac{d[\text{Cr(VI)}(\text{Hydrochloric acid})]}{dt} = k[\text{Cr(VI)}][\text{Malonic acid}][\text{Tartaric acid}][\text{H}^+]$$

## Conclusion

The results obtained from above investigation shows that the end product of

oxidation of Malonic acid + Tartaric acid is Carbondioxide and Carbonyl di carboxylic acid Which may confirmed by their usual test. Several workers reported that the reaction between Malonic acid and Tartaric acid does not involve chain reaction, same observations are seen by author too. In view of author's observations it can be concluded that the retardation of the rate of oxidation of mixture of Malonic acid +Tartaric acid by Chromic acid in presence of Salt, is due to negative catalytic effect. Release of Carbon dioxide, in the oxidation of Tartaric acid by Chromic acid, shows that Chromic acid is attacking the –COOH group of Tartaric acid, not –COOH group of Malonic acid. Oxidation should therefore; take place on the methylene group of Malonic acid and it should involve two electron step. Hence proton of Malonic acid will be eliminated when Chromic acid attacked on  $> \text{CH}_2$  and hence Carbonyl di carboxylic acid *i.e.* Meso Oxalic acid is obtained which is not oxidized by Chromic acid, while Tartaric acid is fully oxidized in to Carbon dioxide.<sup>27-36</sup>

## References

1. A.A. Frost and R.G. Pearson "Kinetics and Mechanism" John Wiley Easter Pvt. Ltd. Publishers P. (150-152), (1970).
2. Ali Mohammad and Leibhafsky J., *Phys. Chem.* 38, 857 (1934).
3. Ansari, A.H. M.P. *Vigyan Academy*, p. 21 (1974).
4. Ansari, A.H., *J., Sci. Res. 1 (i)*, 1-5 (1978) *An. Quim. Ser. A84(II)*, 14-18(1988).
5. Ansari, A.H., Ph.D. Thesis Bhopal Univ. Bhopal (1976).
6. 7 Ansari, A.H., Pramod Kumar & Mishra, B.K. A.K. Singh *J. of Ultra Chem.*, Vol. 4(2), 295-300(2008) *Asian J. of Chemistry*

- 15(3 & 4) 1313-1319 (2003).
8. Bari N.S., Ph.D. Thesis Barkatullah Univ. Bhopal (2002).
  9. Bakore and Shankar *Indian J. Chem. Soc. I (7)*, 286-288 (1963).
  10. Bhagwat and Yadav *J. Ind. Chem. Soc. 41 (1)* 2518 (1964).
  11. Cohen and Westheimer *J. Am. Chem. Soc. 74*, 4287 (1952).
  12. Dharmadhikari Pure, J.E. Ph.D. Thesis Saugar Univ., Sagar (1968).
  13. Das A.K. and Das M., *Ind. J. Chem. 34A*, 866 (1995).
  14. Ekta Pandey, Neeti Grover, Neetu Kamboad & Santosh *Ind. J. Chem. 43 A* 1186-1192 (2004).
  15. Eyring and Eyring "Modern Chemical Kinetics" Reinhold Pub. Corporation London p. 88 (1965).
  16. Friedman and Jafferson, *J. Am. Chem. Soc. 61*, 118 (1939).
  17. Fiegl, F. and Caldas, A., *Anal. Chem. Soc. 35*, 799 (1879).
  18. Fiegl, F., "Spot Tests" Inorganic Analysis Elsevier pub. Comp. Amsterdam, London
  19. (a) Hinshelwood "Kinetics of Chemical Change", Oxford p. 70 & 161 (1949).
  20. Jain Vandana, Ansari, A.H., *J. of Ultra Chem.*, Vol. 10 (1) 70-76 (2014).
  21. Jain Vandana Ph.D. Thesis Barkatullah Univ. Bhopal (2015).
  22. Jain Vandana, Khare Deepti Youth India 2020, unmask the future, 55-57 (2016)
  22. Kabir-ud Din, Khaled Hartani & Zaheer K. *Trans. Met. Chem. 25*, 478-484 (2000).
  23. Mahajani, A.V., Ph.D. Thesis, Saugar Univ. Sagar (1962).
  24. Laidler K. J. "Chemical Kinetics" International Chemical Series, p. 12 (1952).
  25. Sengupta, Kalyankali, Samantha Tapashi, Basu, Samendra Nath. *Tetrahedron*, 42 (20), 5707-5714 (1986).
  26. Sengupta *J. Ind. Chem. Soc.* 42 (10), 725-727 (1965).
  27. Sharma S.N., Ph.D. Thesis, Saugar Univ. Sagar (1965).
  28. Zaheer Khan and Kabiruddin *Ind. J. of Chem. 40 A*, 528-532 (2001).
  29. D.P. Singh, B.K. Mishra and A.H. Ansari *Ultra J. of Chem. Vol. 3(2)* (2007).
  30. D.P. Singh, B.K. Mishra and Khare Rajesh *Ultra J. of Chem. Vol. 6(2)*, (2010)
  31. Westheimer, F.H., *J. Chem. Phys.*, 17, 61, (1949).
  32. Waters "Chemistry of Free Radicals", Oxford. P. 17 (1948).
  33. Ali S. Haq, Kabir-ud-din Dowson M.W. Maddoc p. (2001).
  34. A.A.P. Khan, A. Momd. S. Bano Inds. & engg. ACS Publ. (2011).
  35. Y. Mohammed and O.A. Komolafe *Oriental J. of Chem. Vol. 26(2)* (2010).
  36. M. Mali, S.N. Khodaskar and N.T. Patel *Asian J. Chem. Vol. 16*, 811-817(2004).